

SCHRÖDINGER FUNCTIONAL AND QUANTIZATION OF GAUGE THEORIES IN THE TEMPORAL GAUGE

G. C. ROSSI

Dipartimento di Fisica, Università di Roma "Tor Vergata"

INFN - Sezione di Roma 2

Via della Ricerca Scientifica - 00133 Rome - ITALY

E-mail: rossig@roma2.infn.it

In the language of Feynman path integrals the quantization of gauge theories is most easily carried out with the help of the Schrödinger Functional (SF). Within this formalism the essentially unique gauge fixing condition is $A_0 = 0$ (temporal gauge), as any other rotationally invariant gauge choice can be shown to be functionally equivalent to the former. In the temporal gauge Gauss' law is automatically implemented as a constraint on the states. States not annihilated by the Gauss operator describe the situation in which external (infinitely heavy) colour sources interact with the gauge field. The SF in the presence of an arbitrary distribution of external colour sources can be expressed in an elegant and concise way.

1. — In the framework of the Feynman path integrals¹ a key role is played by the Schrödinger Functional (SF), $K(\eta_2, T_2; \eta_1, T_1) = K(\eta_2, \eta_1; T_2 - T_1)$, which represents the probability amplitude of finding the fundamental fields of the theory, $\{\eta(\mathbf{x}, t)\}$, in the configuration $\eta_2(\mathbf{x})$ at time T_2 , if they were in the configuration $\eta_1(\mathbf{x})$ at time T_1 . $K(\eta_2, \eta_1; T_2 - T_1)$ is the matrix element of the time evolution operator between eigenstates of the field operator, in the Schrödinger representation². In formulae we would schematically write ($T = T_2 - T_1$)

$$K(\eta_2, \eta_1; T) = \langle \eta_2 | e^{-i\mathcal{H}T/\hbar} | \eta_1 \rangle$$

$$\hat{\eta}(\mathbf{x}, t_o) | \eta_I \rangle = \eta_I(\mathbf{x}) | \eta_I \rangle \quad I = 1, 2 \quad (1)$$

where the “hat” on $\hat{\eta}(\mathbf{x}, t_o)$ has been introduced to make clear in the last equation the distinction between field operators and their corresponding eigenvalues, $\eta_I(\mathbf{x})$. t_o is a fixed, arbitrary time, at which the Schrödinger representation is defined. The knowledge of $K(\eta_2, \eta_1; T)$ is sufficient to compute any physical quantity, as seen from the formal expansion (spectral representation)

$$K(\eta_2, \eta_1; T) = \langle \eta_2 | e^{-i\mathcal{H}T/\hbar} | \eta_1 \rangle = \sum_k e^{-iE_k T/\hbar} \Psi_k(\eta_2) \Psi_k^*(\eta_1) \quad (2)$$

where the state functionals $\Psi_k(\eta) = \langle \eta | \Psi_k \rangle$ are eigenstates of the Hamiltonian, $\mathcal{H}$, of the system, $\mathcal{H} | \Psi_k \rangle = E_k | \Psi_k \rangle$.

In this talk I wish to illustrate the quantization of a gauge theory in the gauge $A_o = 0$, using the notion of SF³. For lack of space I will concentrate on the case of pure Yang-Mills (YM) theories and limit the discussion to the topologically trivial sector of the theory. The procedure for including fermions is not entirely trivial and can be found in⁴. Topology is discussed in the third paper of ref.³. For applications of the notion of SF to lattice QCD see⁵.

2. – The elegance of the principle of gauge invariance is counterbalanced by the necessity of introducing the non-canonical variable A_o . In computing physical quantities one is then confronted with the problem of eliminating it by the use of some gauge fixing condition. The simplest choice is to set $A_o = 0$. For obvious reasons this gauge is often referred to as the temporal gauge⁶.

Some nice features of the temporal gauge are the absence of ghosts, the fact that A^i and $\Pi^i = \dot{A}^i$ ($i = 1, 2, 3$) are canonically conjugate variables and the natural interpretation of the solutions of the Euclidean field equations (instantons) as configurations interpolating between vacua (pure gauges) with winding number differing by one unit (tunneling). In the $A_o = 0$ gauge, however, Gauss' law is apparently lost. I will show that Gauss' law is, in fact, naturally implemented as a constraint on the states³.

In the language of path integrals the SF of a YM theory, $K_{YM}(\mathbf{A}_2, \mathbf{A}_1; T)$, can be represented by the formal expression

$$K_{YM}(\mathbf{A}_2, \mathbf{A}_1; T) \sim \int_{\mathbf{A}_1(\mathbf{x})}^{\mathbf{A}_2(\mathbf{x})} \prod_x [d\mathbf{A}(x) dA_o(x)] \exp(i \int_{T_1}^{T_2} \mathcal{L}_{YM}[A_\mu] dx dt) \quad (3)$$

where the functional integration is carried out over all field configurations such that $\mathbf{A}(\mathbf{x}, T_{1,2}) = \mathbf{A}_{1,2}(\mathbf{x})$. $\mathcal{L}_{YM}[A_\mu] = -\frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu}$ is the YM Lagrangian and $SU(N_c)$ is the gauge group. As A_o is the Lagrange multiplier enforcing Gauss' law, the integration over A_o in (3) is extended to all times $T_1 \leq t \leq T_2$ (including the boundaries).

Of course the integration in (3) yields an infinite result as the integrand is invariant under local gauge transformations (g_o is the YM coupling constant)

$$\begin{aligned} A_\mu(x) &\rightarrow A_\mu^{U_h}(x) = U_h^\dagger(x) A_\mu(x) U_h(x) + \frac{i}{g_o} U_h^\dagger(x) \partial_\mu U_h(x) \\ U_h(x) &= \exp[ig_o h(x)] \\ A_\mu(x) - A_\mu^{U_h}(x) &\simeq D_\mu(A) h(x) = \partial_\mu h(x) - ig_o [A_\mu(x), h(x)] \end{aligned} \quad (4)$$

To give a meaning to eq. (3) we introduce in the path integral the identity

$$1 = \Delta_{FP}[A_o] \int_{\mathcal{G}} \prod_{\mathbf{x}} \prod_{t \in [T_1, T_2]} \mathcal{D}h(\mathbf{x}, t) \delta(A_o^{U_h}) \quad (5)$$

where $\mathcal{G}$ is the group of topologically trivial gauge transformations and by $\prod \mathcal{D}h$ we mean the product of the Haar measures of the gauge group $SU(N_c)$ over all space-time points. $\Delta_{FP}[A_o]$ is the usual Fadeev-Popov (FP) factor, which in this gauge only depends on A_o . From eqs. (4) and (5) one obtains

$$K_{YM}(\mathbf{A}_2, \mathbf{A}_1; T) \sim \int \prod_{\mathbf{x}} \prod_{t \in [T_1, T_2]} \mathcal{D}h(\mathbf{x}, t) \cdot \int_{\mathbf{A}_1(\mathbf{x})}^{\mathbf{A}_2(\mathbf{x})} \prod_x [d\mathbf{A}(x) dA_o(x)] \Delta_{FP}[A_o] \delta(A_o^{U_h}) \exp(i \int_{T_1}^{T_2} \mathcal{L}_{YM}[A_\mu] d\mathbf{x} dt) \quad (6)$$

Putting $A_\mu^{U_h} = A'_\mu$, the A'_o integration is immediately carried out with the help of the δ -function. Dropping the field independent FP factor, one gets

$$K_{YM}(\mathbf{A}_2, \mathbf{A}_1; T) \sim \left[\int \prod_{\mathbf{x}} \prod_{t \in (T_1, T_2)} \mathcal{D}h(\mathbf{x}, t) \right] \int_{\mathcal{G}_o} \mathcal{D}h_2(\mathbf{x}) \int_{\mathcal{G}_o} \mathcal{D}h_1(\mathbf{x}) \cdot \int_{\mathbf{A}_1(\mathbf{x})}^{\mathbf{A}_2(\mathbf{x})} \prod_x d\mathbf{A}(x) \exp(i \int_{T_1}^{T_2} \mathcal{L}_{YM}[\mathbf{A}, A_o = 0] d\mathbf{x} dt) \quad (7)$$

where we have separated out the gauge integrations at the initial and final time and introduced the definitions $U_h(\mathbf{x}, T_I) = U_{h_I}(\mathbf{x})$, $I = 1, 2$. The $U_{h_{1,2}}$ integrations are extended over the group, $\mathcal{G}_o$, of the (topologically trivial) time independent gauge transformations. As the integrand does not depend on the gauge transformations at times $t \in (T_1, T_2)$, one can drop the infinite gauge volume, $\mathcal{V} = \int \prod_{\mathbf{x}} \prod_{t \in (T_1, T_2)} \mathcal{D}h(\mathbf{x}, t)$, from the previous equation and write

$$K_{YM}(\mathbf{A}_2, \mathbf{A}_1; T) = \int_{\mathcal{G}_o} \mathcal{D}h_1(\mathbf{x}) \int_{\mathcal{G}_o} \mathcal{D}h_2(\mathbf{x}) \tilde{K}_{YM}(\mathbf{A}_2^{U_{h_2}}, \mathbf{A}_1^{U_{h_1}}; T) \quad (8)$$

$$\tilde{K}_{YM}(\mathbf{A}_2, \mathbf{A}_1; T) = \int_{\mathbf{A}_1(\mathbf{x})}^{\mathbf{A}_2(\mathbf{x})} \prod_x d\mathbf{A}(x) \exp(i \int_{T_1}^{T_2} \mathcal{L}_{YM}[\mathbf{A}, A_o = 0] d\mathbf{x} dt)$$

It should be immediately noted that, as a consequence of the invariance $\tilde{K}_{YM}(\mathbf{A}_2^V, \mathbf{A}_1^V; T) = \tilde{K}_{YM}(\mathbf{A}_2, \mathbf{A}_1; T)$, $V \in \mathcal{G}_o$, the integrand in the first of eqs. (8) only depends on $U_{h_2} U_{h_1}^\dagger$. Consequently one of the two gauge integrations must be dropped in order to get a finite answer⁷, leading to the final formula

$$K_{YM}^{A_o}(\mathbf{A}_2, \mathbf{A}_1; T) = \int_{\mathcal{G}_o} \mathcal{D}h(\mathbf{x}) \tilde{K}_{YM}(\mathbf{A}_2^{U_h}, \mathbf{A}_1; T) \quad (9)$$

From this equation one can construct a perfectly well defined perturbative expansion with no singularity whatsoever in the gluon propagator³ and prove that no problems arise with the time exponentiation of the Wilson loop⁸.

To understand the physical meaning of the gauge “projections” in the above formulae it is more convenient to refer our considerations to eqs. (8). Inserting the spectral decomposition of $\tilde{K}_{YM}(\mathbf{A}_2, \mathbf{A}_1; T)$,

$$\tilde{K}_{YM}(\mathbf{A}_2, \mathbf{A}_1; T) = \langle \mathbf{A}_2 | e^{-i\mathcal{H}_{YM}T/\hbar} | \mathbf{A}_1 \rangle = \sum_{\gamma} e^{-iE_{\gamma}T/\hbar} \Phi_{\gamma}(\mathbf{A}_2) \Phi_{\gamma}^*(\mathbf{A}_1) \quad (10)$$

$$\mathcal{H}_{YM} \Phi_\gamma(\mathbf{A}) = E_\gamma \Phi_\gamma(\mathbf{A}) \quad \mathcal{H}_{YM} = \frac{1}{2} \int d\mathbf{x} \left[-\frac{\delta^2}{\delta A_i^a \delta A_i^a} + \frac{1}{2} F_{ij}^a F_{ij}^a \right] \quad (11)$$

in (8), we get

$$\begin{aligned} K_{YM}(\mathbf{A}_2, \mathbf{A}_1; T) &= \sum_\gamma e^{-iE_\gamma T/\hbar} \int_{G_o} \mathcal{D}h_2 \Phi_\gamma(\mathbf{A}_2^{U_{h_2}}) \int_{G_o} \mathcal{D}h_1 \Phi_\gamma^*(\mathbf{A}_1^{U_{h_1}}) \equiv \\ &\equiv \sum_k e^{-iE_k T/\hbar} \Psi_k(\mathbf{A}_2) \Psi_k^*(\mathbf{A}_1) \implies \Psi_k(\mathbf{A}) = \int_{G_o} \mathcal{D}h \Phi_k(\mathbf{A}^{U_h}) \end{aligned} \quad (12)$$

We see from eqs. (12) that the states appearing in K_{YM} are gauge “projections” of the states in $\tilde{K}_{YM}$, hence they are invariant under G_o . In fact from known properties of the Haar measure, one immediately proves $\Psi_k(\mathbf{A}^V) = \Psi_k(\mathbf{A})$. This means that the functionals Ψ_k in (12) are physical states in the sense that, besides being eigenstates of $\mathcal{H}$, they also satisfy Gauss’ law:

$$0 = \frac{\delta \Psi_k(\mathbf{A}^{U_h})}{\delta h^a(\mathbf{x})} \Big|_{h=0} = \int d\mathbf{y} \frac{\delta \Psi_k(\mathbf{A})}{\delta A_i^b(\mathbf{y})} \frac{\delta (A_i^b(\mathbf{y})^{U_h})}{\delta h^a(\mathbf{x})} \Big|_{h=0} = D_i^{ab}(\mathbf{A}) \frac{\delta \Psi_k(\mathbf{A})}{\delta A_i^b(\mathbf{x})} \quad (13)$$

The chain of equalities in eq. (13) shows that the operator $D_i^{ab}(\mathbf{A}) \delta / \delta A_i^b(\mathbf{x})$ is (up to a factor) the generator of time independent gauge transformations.

It is important to note at this point that the scalar product of states obeying Gauss’ law must be defined by dividing out the volume of the group G_o (as was done in going from eqs. (8) to (9))⁷. A practical way of doing it is to resort to the FP trick and write

$$(\Psi_1, \Psi_2) = \int d\mathbf{A} \Psi_1^*(\mathbf{A}) \Psi_2(\mathbf{A}) \Delta_{FP}^{\mathcal{F}}[\mathbf{A}] \delta(\mathcal{F}(\mathbf{A})) \quad (14)$$

where $\mathcal{F}(\mathbf{A}) = 0$ is any (rotationally invariant) spatial gauge condition (e.g. $\nabla \mathbf{A} = 0$) and $\Delta_{FP}^{\mathcal{F}}[\mathbf{A}]$ is the corresponding FP factor.

3. – We now want to prove the equivalence of the temporal gauge with any other (rotationally invariant) spatial gauge. To be precise we want to show that the SF in the gauge $\mathcal{F}(\mathbf{A}) = 0$, call it $K_{YM}^{\mathcal{F}}$, is equal to the SF, K_{YM} , given by eqs. (10), when the latter is restricted to boundary fields satisfying the gauge conditions $\mathcal{F}(\mathbf{A}_{1,2}) = 0$.

To be concrete let us consider the Coulomb case, in which $\mathcal{F}(\mathbf{A}) = \nabla \mathbf{A}$. As it will be clear, however, our proof is valid for any (rotationally invariant) gauge fixing condition. In the Coulomb gauge the SF, K_{YM}^C , takes the form

$$\begin{aligned} K_{YM}^C(\mathbf{A}_2, \mathbf{A}_1; T) &= \int_{\mathbf{A}_1}^{\mathbf{A}_2} \prod_x dA_\mu(x) \Delta_{FP}^C[\mathbf{A}] \delta(\nabla \mathbf{A}) \exp(i \int_{T_1}^{T_2} \mathcal{L}_{YM}[A_\mu] d\mathbf{x} dt) \\ 1 &= \Delta_{FP}^C[\mathbf{A}] \int_G \prod_{\mathbf{x}} \prod_{t \in (T_1, T_2)} \mathcal{D}h(\mathbf{x}, t) \delta(\nabla \mathbf{A}^{U_h}) \end{aligned} \quad (15)$$

where the second equation defines the FP factor, Δ_{FP}^C . No functional gauge integrations at $t = T_2$ and $t = T_1$ appear in the above formulae as in the Coulomb gauge the boundary fields necessarily obey the gauge conditions $\nabla \mathbf{A}_{1,2} = 0$.

We now perform the change of variables $(A_o, \mathbf{A}) \rightarrow (h, \mathbf{A}')$ defined by

$$\begin{aligned} \mathbf{A}'(\mathbf{x}, t) &= \mathbf{A}^{U_h}(\mathbf{x}, t) = U_h^\dagger(\mathbf{x}, t) \mathbf{A}(\mathbf{x}, t) U_h(\mathbf{x}, t) + \frac{i}{g_o} U_h^\dagger(\mathbf{x}, t) \nabla U_h(\mathbf{x}, t) \\ A'_o(\mathbf{x}, t) &\equiv U_h^\dagger(\mathbf{x}, t) A_o(\mathbf{x}, t) U_h(\mathbf{x}, t) + \frac{i}{g_o} U_h^\dagger(\mathbf{x}, t) \partial_o U_h(\mathbf{x}, t) = 0 \end{aligned} \quad (16)$$

for all $t \in [T_1, T_2]$. Solving the second of eqs. (16) for U_h , one gets $U_h(\mathbf{x}, t) = T \exp(i g_o \int_{T_1}^t A_o(\mathbf{x}, \tau) d\tau) U_h(\mathbf{x}, T_1)$, from which one computes the integration measure in the new variables to be $\delta \mathbf{A} \delta A_o = \delta \mathbf{A}' J(h) \delta h = \delta \mathbf{A}' \mathcal{D}h$. The appearance of the Haar measure in the last equality follows upon observing that

$$J(h) = \det \frac{\delta A_o^a}{\delta h^b} = \det \frac{e^{i\chi} - 1}{i\chi}, \quad \chi_{ab} = i f^{abc} h^c \quad (17)$$

Since from eqs. (16) one gets $\mathcal{L}_{YM}[A_\mu] = \mathcal{L}_{YM}[A_\mu^{U_h}] = \mathcal{L}_{YM}[\mathbf{A}', A'_o = 0]$, K_{YM}^C (see eqs. (15)) can be written in the form

$$\begin{aligned} K_{YM}^C(\mathbf{A}_2, \mathbf{A}_1; T) &= \int \prod_{\mathbf{x}} \prod_{t \in [T_1, T_2]} \mathcal{D}h(\mathbf{x}, t) \int_{\mathbf{A}_1^{U_h(\mathbf{x}, T_1)}}^{\mathbf{A}_2^{U_h(\mathbf{x}, T_2)}} \cdot \\ &\cdot \prod_x d\mathbf{A}'(x) \Delta_{FP}^C[\mathbf{A}'] \delta(\nabla \mathbf{A}'^{U_h^\dagger}) \exp(i \int_{T_1}^{T_2} \mathcal{L}_{YM}[\mathbf{A}', A'_o = 0] dx dt) \end{aligned} \quad (18)$$

Exploiting backwards the definition of Δ_{FP}^C given in eq. (15), we can trivially perform the gauge integrations for all $T_1 < t < T_2$. In this way the whole Coulomb gauge fixing term gets replaced by unity. Only the initial and final gauge integrations are left out: it is then immediately seen that formula (18) coincides with the expression given by eqs. (8), as announced.

4. – External (infinitely heavy) colour sources can be elegantly introduced in the present formalism by noticing that from the invariance property of $\tilde{K}_{YM}$

$$\sum_{\gamma} e^{-iE_{\gamma}T/\hbar} \Phi_{\gamma}(\mathbf{A}_2^V) \Phi_{\gamma}^*(\mathbf{A}_1^V) = \sum_{\gamma} e^{-iE_{\gamma}T/\hbar} \Phi_{\gamma}(\mathbf{A}_2) \Phi_{\gamma}^*(\mathbf{A}_1), \quad V \in \mathcal{G}_o \quad (19)$$

it follows that either **i)** Φ_{γ} is not degenerate and consequently $\Phi_{\gamma}(\mathbf{A}^V) = \Phi_{\gamma}(\mathbf{A})$, implying that Φ_{γ} is actually a physical state annihilated by Gauss' law, or **ii)** there is some degeneracy and we have $(\gamma = n, s) \Phi_{n,s}(\mathbf{A}^V) = R_s^{s'}(V) \Phi_{n,s'}(\mathbf{A})$, where $R(V)$ is some unitary representation of the group $\mathcal{G}_o$.

It has been proved in the first paper of ref. ³ that all unitary finite dimensional representations of $\mathcal{G}_o$ are of the form $R(V_w) = \otimes_{j=1}^p \exp[i g_o w^a(\mathbf{y}_j) M_{R_j}^a]$, where the $M_{R_j}^a$'s are the generators of $SU(N_c)$ in the representation R_j .

To understand the physical meaning of the situation described in **ii)** let us consider the simple case in which $R(V_w) = \exp[ig_\circ w^a(\mathbf{y})M_R^a]$. Then a state transforming according to the representation R , $\Phi_{n,s}^R$, will obey the equation

$$D_i^{ab}(\mathbf{A}) \frac{\delta \Phi_{n,s}^R(\mathbf{A})}{\delta A_i^b(\mathbf{x})} = -ig_\circ [M_R^a]_s^{s'} \Phi_{n,s'}^R(\mathbf{A}) \delta(\mathbf{x} - \mathbf{y}) \quad (20)$$

Eq. (20) is telling us that $\Phi_{n,s}^R$ is not annihilated by the Gauss operator, but describes the dynamics of a point-like external (infinitely heavy) colour source, belonging to the representation R and located in space at the point $\mathbf{y}$, in interaction with the gauge field .

From the point of view of this analysis the formulae given in eqs. (8) should be interpreted as yielding the projection of $\tilde{K}_{YM}$ on the trivial representation of $\mathcal{G}_\circ$. This simple observation gives us a way to generalize eqs. (8) to encompass the construction of the SF in the presence of arbitrary external colour sources. To this end it is enough to project $\tilde{K}_{YM}$ on the appropriate (unitary) representation of $\mathcal{G}_\circ$. To illustrate this procedure in a simple, but physically relevant, case we give the formula for the SF when the gauge field is coupled to a pair of $q\bar{q}$ -like sources. If $\mathbf{x}_q$ and $\mathbf{x}_{\bar{q}}$ are the locations of the two sources in space, one finds

$$K_{YM}^{q\bar{q}}(\mathbf{A}_2, s_2, r_2; \mathbf{A}_1, s_1, r_1; T) = \int_{\mathcal{G}_\circ} \mathcal{D}h \tilde{K}_{YM}(\mathbf{A}_2^{U_h}, \mathbf{A}_1; T) [U_h(\mathbf{x}_q)]_{s_2}^{s_1} [U_h^\dagger(\mathbf{x}_{\bar{q}})]_{r_1}^{r_2}$$

where now the state functionals contributing to $K_{YM}^{q\bar{q}}$ have two extra indices, $s, r = 1, \dots, N_c$, corresponding to the colour degrees of freedom of the sources.

References

1. R.P. Feynman and A.R. Hibbs, *Quantum Mechanics and Path Integrals* (McGraw-Hill Book Co., New York, 1965).
2. K. Symanzik, *Nucl. Phys. B* **190**, 1 (1981).
3. G.C. Rossi and M. Testa, *Nucl. Phys. B* **163**, 109 (1980); *Nucl. Phys. B* **176**, 477 (1980); *Nucl. Phys. B* **237**, 442 (1984).
4. G.C. Rossi and K. Yoshida, *Nuovo Cimento* **11**, 101 (1989); J.P. Leroy, J. Micheli, G.C. Rossi and K. Yoshida, *Z. Phys. C* **48**, 653 (1990).
5. M. Lüscher, Les Houches Lectures, 1997 (hep-lat/9802029) and references therein; S. Sint *Nucl. Phys. B* **421**, 135 (1994).
6. H. Weyl, *The theory of groups and quantum mechanics* (Dover, 1950).
7. G.C. Rossi and M. Testa, *Phys. Rev. D* **29**, 2997 (1984).
8. S. Caracciolo, G. Curci and P. Menotti, *Phys. Lett. B* **113**, 311 (1982).